

“JUST THE MATHS”

SLIDES NUMBER

2.4

SERIES 4

(Further convergence and divergence)

by

A.J.Hobson

<p>2.4.1 Series of positive and negative terms 2.4.2 Absolute and conditional convergence 2.4.3 Tests for absolute convergence 2.4.4 Power series</p>

UNIT 2.4 - SERIES 4 - FURTHER CONVERGENCE AND DIVERGENCE

2.4.1 SERIES OF POSITIVE AND NEGATIVE TERMS

TEST 1 - The r -th Term Test (Revisited)

The r -th Term Test in Unit 2.3 may be used for series whose terms are not necessarily all positive.

Outline Proof:

The formula

$$u_r = S_r - S_{r-1}$$

is valid for any series.

The series cannot converge unless the partial sums S_r and S_{r-1} both tend to the same finite limit as r tends to infinity.

Hence, u_r tends to zero as r tends to infinity.

Alternating Series

A simple kind of series with both positive and negative terms is one whose terms are alternately positive and negative.

Test 4 - The Alternating Series Test

If

$$u_1 - u_2 + u_3 - u_4 + \dots, \text{ where } u_r > 0,$$

is such that

$$u_r > u_{r+1} \text{ and } u_r \rightarrow 0 \text{ as } r \rightarrow \infty,$$

then the series converges.

Outline Proof:

(a) Re-group the series as

$$(u_1 - u_2) + (u_3 - u_4) + (u_5 - u_6) + \dots;$$

That is,

$$\sum_{r=1}^{\infty} v_r$$

where $v_1 = u_1 - u_2, v_2 = u_3 - u_4, v_3 = u_5 - u_6, \dots$

v_r is positive, so that

$$S_r = v_1 + v_2 + v_3 + \dots + v_r$$

increases as r increases.

(b) Alternatively, re-group the series as

$$u_1 - (u_2 - u_3) - (u_4 - u_5) - (u_6 - u_7) - \dots ;$$

That is,

$$u_1 - \sum_{r=1}^{\infty} w_r$$

where $w_1 = u_2 - u_3, w_2 = u_4 - u_5, w_3 = u_6 - u_7, \dots$

$S_r = u_1 - (w_1 + w_2 + w_3 + \dots + w_r)$ is less than u_1 since positive quantities are being subtracted from it.

(c) We conclude that the partial sums of the original series are steadily increasing but are never greater than u_1 .

They must therefore tend to a finite limit as r tends to infinity; that is, the series converges.

ILLUSTRATION

The series

$$\sum_{r=1}^{\infty} (-1)^{r-1} \frac{1}{r} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

is convergent since

$$\frac{1}{r} > \frac{1}{r+1} \quad \text{and} \quad \frac{1}{r} \rightarrow 0 \quad \text{as} \quad r \rightarrow \infty.$$

2.4.2 ABSOLUTE AND CONDITIONAL CONVERGENCE

DEFINITION (A)

If

$$\sum_{r=1}^{\infty} u_r$$

is a series with both positive and negative terms, it is said to be “**absolutely convergent**” if

$$\sum_{r=1}^{\infty} |u_r|$$

is convergent.

DEFINITION (B)

If

$$\sum_{r=1}^{\infty} u_r$$

is a convergent series of positive and negative terms, but

$$\sum_{r=1}^{\infty} |u_r|$$

is a divergent series, then the first of these two series is said to be “**conditionally convergent**”.

ILLUSTRATIONS

1. The series

$$1 + \frac{1}{2^2} - \frac{1}{3^2} - \frac{1}{4^2} + \frac{1}{5^2} + \frac{1}{6^2} + \dots$$

converges absolutely since the series

$$1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \frac{1}{6^2} + \dots$$

converges.

2. The series

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots$$

is conditionally convergent.

It converges (by the Alternating Series Test), but the series

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots \text{ diverges.}$$

Notes:

(i) It may be shown that any series of positive and negative terms which is **absolutely** convergent will also be convergent.

(ii) Any test for the convergence of a series of positive terms may be used as a test for the absolute convergence of a series of both positive and negative terms.

2.4.3 TESTS FOR ABSOLUTE CONVERGENCE

The Comparison Test

Given the series

$$\sum_{r=1}^{\infty} u_r,$$

suppose that $|u_r| \leq v_r$ where

$$\sum_{r=1}^{\infty} v_r$$

is a convergent series of positive terms. Then, the given series is absolutely convergent.

D'Alembert's Ratio Test

Given the series

$$\sum_{r=1}^{\infty} u_r,$$

suppose that

$$\lim_{r \rightarrow \infty} \left| \frac{u_{r+1}}{u_r} \right| = L.$$

Then the given series is absolutely convergent if $L < 1$.

Note:

If $L > 1$, then $|u_{r+1}| > |u_r|$ for large enough values of r showing that the **numerical** values of the terms steadily increase.

This implies that u_r does **not** tend to zero as r tends to infinity

Hence, (by the r -th Term Test) the series diverges.

If $L = 1$, there is no conclusion.

EXAMPLES

1. Show that the series

$$\frac{1}{1 \times 2} - \frac{1}{2 \times 3} - \frac{1}{3 \times 4} + \frac{1}{4 \times 5} - \frac{1}{5 \times 6} - \frac{1}{6 \times 7} + \dots$$

is absolutely convergent.

Solution

The r -th term of the series is numerically equal to

$$\frac{1}{r(r+1)}$$

This is always less than $\frac{1}{r^2}$, the r -th term of a known convergent series.

2. Show that the series

$$\frac{1}{2} - \frac{2}{5} + \frac{3}{10} - \frac{4}{17} + \dots$$

is conditionally convergent.

Solution

The r -th term of the series is numerically equal to

$$\frac{r}{r^2 + 1},$$

which tends to zero as r tends to infinity.

Also,

$$\frac{r}{r^2 + 1} > \frac{r + 1}{(r + 1)^2 + 1}$$

since this may be reduced to the true statement $r^2 + r > 1$.

Hence, by the alternating series test, the series converges.

However,

$$\frac{r}{r^2 + 1} > \frac{r}{r^2 + r} = \frac{1}{r + 1}$$

and, hence, by the Comparison Test, the series of absolute values is divergent since

$$\sum_{r=1}^{\infty} \frac{1}{r + 1}$$

is divergent.

2.4.4 POWER SERIES

A series of the form

$$a_0 + a_1x + a_2x^2 + a_3x^3 + \dots = \sum_{r=0}^{\infty} a_r x^r \quad \text{or} \quad \sum_{r=1}^{\infty} a_{r-1} x^{r-1},$$

where x is usually a variable quantity, is called a “**power series in x with coefficients,**

$a_0, a_1, a_2, a_3, \dots$ ”.

Notes:

(i) By summing the series from $r = 0$ to infinity, the constant term at the beginning (if there is one) can be considered as the term in x^0 .

The various tests for convergence and divergence still apply in this alternative notation.

(ii) A power series will not necessarily be convergent (or divergent) for **all** values of x

It is usually required to determine the specific **range** of values of x for which the series converges

This can most frequently be done using D’Alembert’s Ratio Test

ILLUSTRATION

For the series

$$x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots = \sum_{r=1}^{\infty} (-1)^{r-1} \frac{x^r}{r},$$

$$\left| \frac{u_{r+1}}{u_r} \right| = \left| \frac{(-1)^r x^{r+1}}{r+1} \cdot \frac{r}{(-1)^{r-1} x^r} \right| = \left| \frac{r}{r+1} x \right|,$$

which tends to $|x|$ as r tends to infinity.

Thus, the series converges absolutely when $|x| < 1$ and diverges when $|x| > 1$.

If $x = 1$, we have the series

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

which converges by the Alternating Series Test.

If $x = -1$, we have the series

$$-1 - \frac{1}{2} - \frac{1}{3} - \frac{1}{4} - \dots$$

which diverges.

The **precise** range of convergence for the given series is therefore $-1 < x \leq 1$.